


NASA TECHNICAL NOTE



NASA TN D-6412

2.1

LOAN COPY: RETURN
AFWL (DOGL)
KIRTLAND AFB, N. M



NASA TN D-6412

DEPENDENCE OF INTEGRATED
CROSS SECTIONS ON THE
ROTATIONAL STATES OF SODIUM-23
IN THE REACTION MAGNESIUM-25 (d, α)
SODIUM-23 AT 20.9 MeV

by Joseph R. Priest

*Lewis Research Center
Cleveland, Ohio 44135*

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION • WASHINGTON, D. C. • JULY 1971



0132838

| | | | | | |
|---|--|--|--|---|--|
| 1. Report No. NASA TN D-6412 | | 2. Government Accession No. | | 3. Recipient's | |
| 4. Title and Subtitle DEPENDENCE OF INTEGRATED CROSS SECTIONS ON THE ROTATIONAL STATES OF SODIUM-23 IN THE REACTION MAGNESIUM-25 (d, α) SODIUM-23 AT 20.9 MeV | | | | 5. Report Date July 1971 | |
| | | | | 6. Performing Organization Code | |
| | | | | 8. Performing Organization Report No. E-6264 | |
| | | | | 10. Work Unit No. 129-02 | |
| 7. Author(s) Joseph R. Priest, Miami University, Oxford, Ohio | | | | 11. Contract or Grant No. | |
| 9. Performing Organization Name and Address Lewis Research Center National Aeronautics and Space Administration Cleveland, Ohio 44135 | | | | 13. Type of Report and Period Covered Technical Note | |
| 12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, D.C. 20546 | | | | 14. Sponsoring Agency Code | |
| 15. Supplementary Notes | | | | | |
| 16. Abstract The integrated cross sections for production of $K = 3/2$ states in the $^{25}\text{Mg}(d, \alpha)^{23}\text{Na}$ reaction obey a linear relation of the form $\sigma = a(2I + 1) - b$. A comparison of these results with similar data from the $^{27}\text{Al}(d, \alpha)^{25}\text{Mg}$ reaction suggests that the negative intercept is a consequence of band mixing in the states of ^{23}Na . | | | | | |
| 17. Key Words (Suggested by Author(s)) Differential cross sections for $^{24}\text{Mg}(d, \alpha)^{23}\text{Na}$ Band mixing compared with $^{27}\text{Al}(d, \alpha)^{25}\text{Mg}$ | | | | 18. Distribution Statement Unclassified - unlimited | |
| 19. Security Classif. (of this report) Unclassified | | 20. Security Classif. (of this page) Unclassified | | 21. No. of Pages 13 | |
| | | | | 22. Price* \$3.00 | |

DEPENDENCE OF INTEGRATED CROSS SECTIONS ON THE ROTATIONAL STATES OF SODIUM-23 IN THE REACTION MAGNESIUM-25 (d, α)

SODIUM-23 AT 20.9 MeV

by Joseph R. Priest*

Lewis Research Center

SUMMARY

The angular distributions of differential cross sections corresponding to production of rotational states with rotational quantum number (K) of $3/2$ from the (d, α) reaction on magnesium 25 (^{25}Mg) were measured from 15° to 165° (laboratory system). The mean deuteron energy was 20.9 MeV (laboratory system). The angular distributions are similar and peaked in the forward direction. This is indicative of a direct reaction mechanism. The integrated cross sections obey a simple linear relation (of the form $\sigma = a(2I + 1) - b$). A comparison of these results with similar data from the $^{27}\text{Al}(d, \alpha)^{25}\text{Mg}$ reaction suggests that the negative intercept is a consequence of rotational band mixing in the states of ^{23}Na .

INTRODUCTION

In a study of the $^{27}\text{Al}(d, \alpha)^{25}\text{Mg}$ reaction at 20.9 MeV, we observed that the angular distributions of differential cross sections for production of states in a given rotational band ($K = 5/2$ and $K = 1/2$) were strikingly similar both in magnitude and shape (ref. 1). The integrated cross sections for the first three states of the $K = 5/2$ and $K = 1/2$ rotational bands were 459, 475, and 417 microbarns and 38, 38, and 95 microbarns, respectively. The noticeably higher value of 95 μb for the $K = 1/2$, $I = 5/2$ state was thought to reflect a mixing of the $I = 5/2$ states of the $K = 5/2$ and $K = 1/2$ rotational bands. This interpretation is consistent with that suggested from inelastic alpha particle scattering experiments on ^{25}Mg (ref. 2). Using the 20.9-MeV results and other data for five deuteron energies between 4.74 and 14.7 MeV, the authors of references 3 and 4

* Professor of Physics, Miami University, Oxford, Ohio.

assumed that the integrated cross section could be decomposed as $\sigma(\alpha_\nu^K) = \sigma_D(\alpha_\nu^K) + \sigma_C(\alpha_\nu^K)$ and showed that for a given energy $\sigma_D(\alpha_\nu^K)$ is nearly constant for each rotational band and that $\sigma_C = a(2I + 1)$. The terms σ_C and σ_D are the contributions due to compound and direct processes, ν corresponds to the ν^{th} state of ^{25}Mg , and K is the rotational band quantum number. In their analysis, our results for a deuteron energy of 20.9 MeV imply that $\sigma_C < 0$. They also observed that for energies above 8 MeV, the values for $\sigma_D(\alpha_4^{1/2})$ were consistently higher than those for $\sigma_D(\alpha_1^{1/2})$ and $\sigma_D(\alpha_2^{1/2})$.

Similar measurements on the $^{25}\text{Mg}(d, \alpha)^{23}\text{Na}$ reaction provide further tests of the systematics observed in the $^{27}\text{Al}(d, \alpha)^{25}\text{Mg}$ reaction, since ^{23}Na is also commonly depicted as a collective-type nucleus. The ground, 0.440-, 2.078-, and 2.705-MeV states (fig. 1) are presumed to be the $I = 3/2^+$, $5/2^+$, $7/2^+$, and $9/2^+$ members of a $K = 3/2$ rotational band. Theoretical calculations (ref. 5) indicate that these states contain admixtures of other states of the same I in rotational bands with $\Delta K = \pm 1$. If the integrated cross sections are sensitive to this band mixing, as they apparently are in the $^{27}\text{Al}(d, \alpha)^{25}\text{Mg}$ reaction, then measurements on the $^{25}\text{Mg}(d, \alpha)^{23}\text{Na}$ reaction should reflect this.

SYMBOLS

| | |
|--|---|
| I | total angular momentum quantum number |
| K | value of the projection of the angular momentum I parallel to the nuclear symmetry axis |
| $\sigma(\alpha_\nu^K)$ | integrated cross section for production of the ν^{th} state having rotational quantum number K |
| $\sigma_D(\alpha_\nu^K), \sigma_C(\alpha_\nu^K)$ | contributions to the integrated cross section due to direct and compound processes |

EXPERIMENTAL PROCEDURE

The ^{25}Mg target was a self-supporting, vacuum-evaporated film having an areal density of 1.203 ± 0.050 milligrams per square centimeter. It was bombarded with the 20.9 ± 0.1 -MeV deuteron beam from the Lewis Research Center cyclotron. The alpha-particle detection and discrimination scheme was the same as in previous (d, α) experiments (refs. 6, 7, and 8). A typical alpha-particle spectrum for a laboratory reaction angle of 20° is shown in figure 2. The alpha-particle groups of interest were fitted with skewed gaussian functions to obtain the experimental yields for calculation of the differential cross sections.

RESULTS AND DISCUSSION

Angular distributions of differential cross sections corresponding to production of the first four $K = 3/2$ states of ^{23}Na in the reaction $^{25}\text{Mg}(d, \alpha)^{23}\text{Na}$ at 20.9 MeV are displayed in figure 3. The numerical data are presented in table I. The overall energy resolution was 200 keV. Consequently, the 2.641- and 2.705-MeV states of ^{23}Na were not resolved. The errors shown in figure 3 and table I reflect the statistical uncertainty of the yields and the error associated with fitting the skewed gaussian functions to the data. The absolute uncertainty is assessed at 15 percent.

We note that these angular distributions have the same basic shape over the entire angular range. The cross sections integrated over angular ranges of 15° to 165° (σ) and 95° to 165° (σ_C) are shown as functions of $2I + 1$ in figure 4. We observe, as did the authors of references 3 and 4 for the $^{27}\text{Al}(d, \alpha)^{25}\text{Mg}$ reaction, that σ and σ_C can be adequately characterized by functions of the form $\sigma = a(2I + 1) + b$ and $\sigma_C = a'(2I + 1)$. We have called σ_C the integral of $d\sigma/d\Omega$ from 95° to 165° . In references 3 and 4, σ_C is called the integral of $d\sigma/d\Omega_{\text{isotropic}}$ over the full angular range studied, where $d\sigma/d\Omega_{\text{isotropic}}$ is the minimum value of $d\sigma/d\Omega$. The form of our curve for σ_C against $2I + 1$ would be unaltered if we used this same interpretation. The authors of references 3 and 4 logically interpreted b as the contribution to the total integrated cross section due to direct processes. In the present study, such an interpretation is not possible because b is clearly negative. A consistent way of interpreting this result would be to say that the admixture in the $K = 3/2$ states from states in the $K = 5/2$ rotational band increases as I increases. This would tend to increase σ as I increases. The theoretical calculations of Kelson and Levinson (ref. 5), which characterize the $3/2^+$, $5/2^+$, and $7/2^+$ states of ^{23}Na with $K = 3/2$ amplitudes of 0.992, 0.862, and 0.777, and $K = 5/2$ amplitudes of 0, 0.453, and 0.556, respectively, would tend to corroborate this reasoning. Additional evidence for these admixtures may be seen in figure 5, where angular distributions for $K = 5/2$ to $K = 5/2$ and $K = 5/2$ to $K = 3/2$ transitions in the reactions $^{27}\text{Al}(d, \alpha)^{25}\text{Mg}$ and $^{25}\text{Mg}(d, \alpha)^{23}\text{Na}$ are displayed. The striking overall similarity both in magnitude and shape indicate that the major admixture in the $K = 3/2$, $I = 9/2$ state of ^{23}Na comes from the $K = 5/2$ rotational band.

CONCLUDING REMARKS

We conclude, that the results of this investigation agree with those from a study of the $^{27}\text{Al}(d, \alpha)^{25}\text{Mg}$ reaction if we account for band mixing in the states of ^{23}Na .

Lewis Research Center,
National Aeronautics and Space Administration,
Cleveland, Ohio, April 27, 1971,
129-02.

REFERENCES

1. Priest, Joseph R. ; and Vincent, John S. : Study of the Reaction $^{27}\text{Al}(d, \alpha)^{25}\text{Mg}$ at 20.9 MeV. Phys. Rev., vol. 152, no. 3, Dec. 16, 1966, pp. 989-994.
2. Blair, J. S. : Inelastic Scattering. Nuclear Spectroscopy with Direct Reactions. II. Proceedings. Rep. ANL-6878, Argonne National Lab., Mar. 1964, pp. 143-173.
3. Hofmann, A. ; Emmerich, W. ; Dinter, H. ; and Philipp, G. : Abhängigkeit des Wirkungsquerschnitts für $^{27}\text{Al}(d, \alpha)^{25}\text{Mg}$ von den Rotationszuständen des ^{25}Mg . Phys. Letters, vol. 26B, no. 6, Feb. 19, 1968, pp. 378-379.
4. Hofmann, A. ; Emmerich, W. ; Dinter, H. ; and Philipp, G. : Die Reaktion $^{27}\text{Al}(d, \alpha)^{25}\text{Mg}$. Nucl. Phys., vol. A113, 1968, pp. 86-96.
5. Kelson, I. ; and Levinson, C. A. : Application of Self-Consistent Field Methods to Rotational Motion in sd Shell Nuclei. Phys. Rev., vol. 134, no. 2B, Apr. 27, 1964, pp. 269-278.
6. Priest, Joseph R. ; and Vincent, John S. : Investigation of the (d, α) Reaction on Aluminum 27. NASA TN D-3548, 1966.
7. Priest, Joseph R. ; and Vincent, John S. : Interpretation of (d, α) Reactions on Fluorine 19 and Nitrogen 15. NASA TN D-3813, 1967.
8. Priest, Joseph R. ; and Vincent, John S. : Differential Cross Sections for (d, α) Reaction on Scandium 45. NASA TN D-4140, 1967.
9. Dubois, J. : A Nuclear Structure Study of ^{23}Na by Means of the $^{22}\text{Ne}(^3\text{He}, d)^{23}\text{Na}$ Reaction. Nucl. Phys., vol. A104, 1967, pp. 657-676.
10. Buechner, W. W. ; and Sperduto, A. : Energy Levels in ^{23}Na and ^{20}Ne . Phys. Rev., vol. 106, no. 5, June 1, 1957, pp. 1008-1011.

TABLE I. - EXPERIMENTAL

(a) Ground state

(b) 0.440-MeV excited state

| Center-of-mass reaction angle, θ_{cm} , deg | Differential cross section, $d\sigma/d\Omega$, $\mu\text{b/sr}$ | Estimated error, $\Delta\sigma$, $\mu\text{b/sr}$ | Center-of-mass reaction angle, θ_{cm} , deg | Differential cross section, $d\sigma/d\Omega$, $\mu\text{b/sr}$ | Estimated error, $\Delta\sigma$, $\mu\text{b/sr}$ |
|--|---|---|--|---|---|
| 16.50 | 33.40 | 1.86 | 16.51 | 102.71 | 2.77 |
| 21.98 | 28.43 | 1.78 | 22.00 | 79.12 | 2.91 |
| 27.45 | 19.84 | 1.19 | 27.47 | 60.66 | 2.03 |
| 32.90 | 18.68 | 1.05 | 32.93 | 46.98 | 1.67 |
| 38.33 | 12.03 | 1.26 | 38.36 | 38.04 | 1.97 |
| 43.73 | 13.71 | 1.04 | 43.76 | 44.67 | 1.85 |
| 49.11 | 10.29 | .74 | 49.14 | 39.93 | 1.38 |
| 54.45 | 9.18 | .67 | 54.49 | 36.23 | 1.35 |
| 59.76 | 8.23 | .59 | 59.80 | 26.89 | .96 |
| 65.03 | 6.75 | .66 | 65.07 | 22.61 | 1.15 |
| 70.27 | 6.08 | .44 | 70.31 | 16.34 | .68 |
| 75.46 | 7.61 | .47 | 75.51 | 14.53 | .64 |
| 80.62 | 7.44 | .51 | 80.66 | 12.53 | .64 |
| 85.73 | 7.83 | .33 | 85.77 | 12.46 | .41 |
| 90.79 | 7.19 | .46 | 90.84 | 11.92 | .60 |
| 95.81 | 6.28 | .42 | 95.86 | 8.79 | .49 |
| 100.79 | 7.62 | 1.98 | 100.84 | 12.33 | 1.97 |
| 105.73 | 8.75 | .57 | 105.77 | 12.13 | .59 |
| 110.62 | 9.19 | .88 | 110.66 | 13.55 | .96 |
| 115.46 | 9.16 | .93 | 115.51 | 12.83 | 1.00 |
| 120.27 | 7.57 | 1.04 | 120.31 | 11.66 | 1.16 |
| 125.03 | 6.63 | .79 | 125.07 | 10.51 | .80 |
| 129.76 | 7.12 | 1.02 | 129.80 | 11.24 | 1.07 |
| 134.45 | 8.10 | .88 | 134.49 | 11.60 | .96 |
| 139.11 | 8.55 | 1.08 | 139.14 | 13.40 | 1.20 |
| 143.73 | 9.20 | .59 | 143.76 | 14.93 | .68 |
| 148.33 | 10.54 | 1.01 | 148.36 | 15.93 | 1.16 |
| 152.90 | 9.88 | 1.11 | 152.93 | 19.10 | 1.23 |
| 157.45 | 14.03 | 2.30 | 157.47 | 23.02 | 2.69 |
| 161.98 | 13.00 | 1.53 | 162.00 | 25.25 | 1.85 |
| 166.50 | 16.63 | 2.06 | 166.51 | 25.43 | 2.37 |

DATA FOR $^{25}\text{Mg}(\text{d}, \alpha)^{23}\text{Na}$

(c) 2.078-MeV excited state

(d) 2.705-MeV excited state

| | | | | | |
|--------|--------|------|--------|--------|------|
| 16.56 | 155.54 | 3.24 | 16.58 | 235.38 | 3.93 |
| 22.06 | 138.09 | 5.35 | 22.09 | 203.93 | 6.62 |
| 27.55 | 101.47 | 3.92 | 27.59 | 161.81 | 4.97 |
| 33.02 | 80.53 | 4.00 | 33.06 | 118.00 | 4.97 |
| 38.47 | 62.52 | 2.38 | 38.51 | 84.81 | 2.80 |
| 43.89 | 60.25 | 3.10 | 43.94 | 83.24 | 3.64 |
| 49.28 | 60.07 | 2.94 | 49.33 | 82.05 | 3.44 |
| 54.64 | 51.13 | 1.87 | 54.69 | 81.04 | 2.37 |
| 59.96 | 42.40 | 1.42 | 60.02 | 72.48 | 1.85 |
| 65.24 | 33.18 | 1.39 | 65.31 | 61.81 | 1.90 |
| 70.49 | 29.25 | 1.36 | 70.56 | 48.52 | 1.79 |
| 75.69 | 22.21 | 1.27 | 75.76 | 40.54 | 1.75 |
| 80.85 | 20.23 | .80 | 80.92 | 35.28 | 1.07 |
| 85.96 | 18.85 | 1.15 | 86.04 | 30.11 | 1.48 |
| 91.03 | 16.24 | .95 | 91.11 | 23.30 | 1.17 |
| 96.06 | 14.28 | 1.01 | 96.13 | 22.35 | 1.30 |
| 101.03 | 17.94 | 1.25 | 101.11 | 25.15 | 1.44 |
| 105.96 | 16.67 | .71 | 106.04 | 26.46 | .87 |
| 110.85 | 17.70 | 1.38 | 110.92 | 27.00 | 1.67 |
| 115.69 | 18.43 | 1.60 | 115.76 | 24.36 | 1.83 |
| 120.49 | 17.83 | 1.54 | 120.56 | 23.59 | 1.76 |
| 125.24 | 16.36 | .78 | 125.31 | 23.01 | .92 |
| 129.96 | 18.48 | 1.81 | 130.02 | 28.96 | 2.20 |
| 134.64 | 18.23 | 1.22 | 134.69 | 32.48 | 1.58 |
| 139.28 | 18.13 | 1.53 | 139.33 | 31.48 | 1.98 |
| 143.89 | 20.45 | 1.16 | 143.94 | 29.99 | 1.41 |
| 148.47 | 21.62 | 2.48 | 148.51 | 31.46 | 3.05 |
| 153.02 | 22.94 | 1.33 | 153.06 | 33.24 | 1.54 |
| 157.55 | 27.85 | 2.96 | 157.59 | 39.76 | 3.49 |
| 162.06 | 28.21 | 2.16 | 162.09 | 36.33 | 2.44 |
| 166.56 | 28.20 | 2.30 | 166.58 | 34.26 | 2.54 |




Figure 1. - The low-lying states of ^{23}Na .
 (The spin assignments were taken from
 ref. 9, the energies from ref. 10.)




Figure 2. - Alpha-particle spectrum obtained at 20° for $^{25}\text{Mg}(d,\alpha)^{23}\text{Na}$ reaction at 20.9 MeV. The solid curve is a least-squares fit of skewed Gaussian functions.




Figure 3. - Angular distributions of differential cross sections for production of $K = 3/2$ states of ^{23}Na in the reaction $^{25}\text{Mg}(d, \alpha)^{23}\text{Na}$ at 20.9 MeV.




Figure 4. - Integrated cross sections as functions of $2I + 1$, where I is the spin quantum number of the final state. (The largest value of σ_C is plotted against $2I + 1 = 11$ because the 2.641- and 2.705-MeV states were unresolved.)




Figure 5. - Angular distributions of differential cross sections for $K = 5/2$ to $K = 5/2$ and $K = 5/2$ to $K = 3/2$ transitions for the reactions $^{27}\text{Al}(d, \alpha)^{25}\text{Mg}$ and $^{25}\text{Mg}(d, \alpha)^{23}\text{Na}$.

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION
WASHINGTON, D. C. 20546
OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE \$300

FIRST CLASS MAIL



POSTAGE AND FEES PAID
NATIONAL AERONAUTICS AND
SPACE ADMINISTRATION

10U 001 49 51 3DS 71166 00903
AIR FORCE WEAPONS LABORATORY /WLOL/
KIRTLAND AFB, NEW MEXICO 87117

ATT E. LOU BOWMAN, CHIEF, TECH. LIBRARY

POSTMASTER: If Undeliverable (Section 158
Postal Manual) Do Not Return

"The aeronautical and space activities of the United States shall be conducted so as to contribute . . . to the expansion of human knowledge of phenomena in the atmosphere and space. The Administration shall provide for the widest practicable and appropriate dissemination of information concerning its activities and the results thereof."

—NATIONAL AERONAUTICS AND SPACE ACT OF 1958

NASA SCIENTIFIC AND TECHNICAL PUBLICATIONS

TECHNICAL REPORTS: Scientific and technical information considered important, complete, and a lasting contribution to existing knowledge.

TECHNICAL NOTES: Information less broad in scope but nevertheless of importance as a contribution to existing knowledge.

TECHNICAL MEMORANDUMS: Information receiving limited distribution because of preliminary data, security classification, or other reasons.

CONTRACTOR REPORTS: Scientific and technical information generated under a NASA contract or grant and considered an important contribution to existing knowledge.

TECHNICAL TRANSLATIONS: Information published in a foreign language considered to merit NASA distribution in English.

SPECIAL PUBLICATIONS: Information derived from or of value to NASA activities. Publications include conference proceedings, monographs, data compilations, handbooks, sourcebooks, and special bibliographies.

TECHNOLOGY UTILIZATION PUBLICATIONS: Information on technology used by NASA that may be of particular interest in commercial and other non-aerospace applications. Publications include Tech Briefs, Technology Utilization Reports and Technology Surveys.

Details on the availability of these publications may be obtained from:

SCIENTIFIC AND TECHNICAL INFORMATION OFFICE
NATIONAL AERONAUTICS AND SPACE ADMINISTRATION
Washington, D.C. 20546